

Quark Masses in the Custodial RS model and how to test it

John N Ng

Triumf

U Montreal , 2008

Outline

- Brief introduction of the Randall-Sundrum Model
- Scalar and Gauge fields in the RS model
- Fermions in RS *location ,location, location*
- The need for custodial $SU(2)_R$. Bulk symmetry is $SU(2)_L \times SU(2)_R \times U(1)_{B-L}$
- Quark Mass Matrices : symmetrical or asymmetrical
- Can we test it in $t \rightarrow jets + Z$ and $t \rightarrow Wb$ at the LHC?
 - (A) KK gauge boson
 - (B) KK Fermions mixing
-
- Conclusions

Introduction to the Randall-Sundrum Model

- There are more than 4 dim. Indeed RS assumes 1 + 4 dim with a warp or conformal metric. AdS
- 5D interval is given by

$$ds^2 = G_{AB} dx^A dx^B = e^{-2kr_c|\phi|} \eta_{\mu\nu} dx^\mu dx^\nu - r_c^2 d\phi^2$$

- Two branes are located at $\phi = 0$ (UV) and $\phi = \pi$ (IR).
- Metric is

$$-\pi \leq \phi \leq \pi, \quad \sigma \equiv kr_c|\phi|,$$
$$G_{AB} = \begin{pmatrix} e^{-2\sigma} \eta_{\mu\nu} & 0 \\ 0 & -r_c^2 \end{pmatrix}, \quad G^{AB} = \begin{pmatrix} e^{+2\sigma} \eta^{\mu\nu} & 0 \\ 0 & -\frac{1}{r_c^2} \end{pmatrix}$$

RS model as 5D field theory

- The action is generalized to 5D e.g. the bulk scalar field we have

$$\begin{aligned} S_5 &= \frac{1}{2} \int d^4x \int_{-\pi}^{\pi} d\phi \sqrt{G} (G^{MN} \partial_M \Phi \partial_N \Phi - m^2 \Phi^2) \\ &= \frac{1}{2} \int d^4x \int_{-\pi}^{\pi} r_c d\phi \left[e^{-2\sigma} \eta^{\mu\nu} \partial_\mu \Phi \partial_\nu \Phi + \frac{1}{r_c^2} \Phi \partial_\phi (e^{-4\sigma} \partial_\phi \Phi) - m^2 e^{-4\sigma} \Phi^2 \right] \end{aligned} \quad (1)$$

- Integrate over ϕ to give a 4D effective theory
- Do KK decomposition.

$$\Phi(x, \phi) = \frac{e^\sigma}{\sqrt{r_c}} \sum_n \Phi_n(x) y_n(\phi)$$

One recovers the canonical 4D scalar field $[\Phi_n] = 1$ and the KK eigen-mode $[y_n] = 0$. y_n is normalized by

$$\int_{-\pi}^{\pi} d\phi y_n(\phi) y_m(\phi) = \delta_{mn}$$

RS:Scalars

- The $y_n(\phi)$ satisfies the eigenvalue eqn

$$-\frac{e^\sigma}{r_c^2} \partial_\phi (e^{-4\sigma} \partial_\phi (e^\sigma y_n)) + m^2 e^{-2\sigma} y_n = m_n^2 y_n$$

- The 4D effective action becomes

$$S = \frac{1}{2} \sum_n \int d^4x [\eta^{\mu\nu} \partial_\mu \Phi_n \partial_\nu \Phi_n - m_n^2 \Phi_n^2]$$

- $m_n = 0$ are the zero modes. Identify them as SM fields.
- The solutions are exponentials

$$y_0 = \frac{e^{kr_c\phi}}{N_0} \left[e^{-\nu kr_c\phi} + b_0 e^{+\nu kr_c\phi} \right], \quad (\nu \equiv \sqrt{4 + m^2/k^2})$$

More RS

- After integrating out the extra the dimension the 4D effective action is

$$S = \frac{1}{2} \sum_n \int d^4x [\eta^{\mu\nu} \partial_\mu \Phi_n \partial_\nu \Phi_n - m_n^2 \Phi_n^2]$$

- For $m_n \neq 0$ the solutions are given by Bessel functions of order $\nu = \sqrt{4} + \frac{m^2}{k^2}$

$$y_n(\phi) = \frac{e^\sigma}{N_n} [J_\nu(z_n) + b_n Y_\nu(z_n)]$$

- m_n and b_n are determined by boundary conditions at $\phi = 0, \pi$. The derivatives are continuous.

Gauge Fields in RS

- Take QED as the toy model. The action is

$$S_5 = -\frac{1}{4} \int d^4x \int_{-\pi}^{\pi} d\phi \sqrt{G} F^{MN} F_{MN}$$

- Choose the unitarity gauge $A_4 = 0$ and KK decompose the gauge field

$$A_\mu(x, \phi) = \frac{1}{\sqrt{r_c}} \sum_n A_\mu^n(x) \chi_n(\phi)$$

with a normalization

$$\int_{-\pi}^{\pi} d\phi \chi_n \chi_m = \delta_{mn}$$

- The 4D Lagrangian is

$$\mathcal{L}_4 \supset +\frac{1}{2} \eta^{\mu\nu} \sum_{m,n} A_\mu^n A_\nu^m \int_{-\pi}^{\pi} d\phi \chi_m \partial_\phi \left(\frac{e^{-2\sigma}}{r_c^2} \partial_\phi \chi_n \right)$$

RS Gage Fields II

- The zero mode has a flat profile

$$\chi_0 = \frac{1}{\sqrt{2\pi}}$$

This preserves charge universality

- The solutions for KK excitations are Bessel functions of order unity

$$\chi_n = \frac{m_n e^\sigma}{k N_n} [J_1(z_n) + a_n Y_1(z_n)]$$

- a_n and m_n are determined by b.c.c at the fixed points $\phi = 0, \pi$

Fermions in 5D Bulk

- 5D fermions are 4-component spinors i.e. vector-like fermions

$$\Psi(x^\mu, y) = \begin{pmatrix} \psi_R(x^\mu, y) \\ \psi_L(x^\mu, y) \end{pmatrix}$$

- The Dirac matrices in 5D are $\gamma^M = (\gamma^\mu, i\gamma^5)$
- Project out the L,R chiral states by boundary conditions or orbifold parities ,i.e. how the field transforms under $Z_2 : y \rightarrow -y$
-

$$\begin{pmatrix} \psi_R(x, y) \\ \psi_L(x, y) \end{pmatrix} \longrightarrow \pm \begin{pmatrix} \psi_R(x, -y) \\ -\psi_L(x, -y) \end{pmatrix}$$

Fermions in Warp Space

- 5D action for fermions is

$$\int d^4x d\phi \sqrt{G} E_a^A \bar{\Psi} \gamma^a D_A \Psi$$

where $E_a^A = \text{diag} \left(e^\sigma, e^\sigma, e^\sigma, e^\sigma, \frac{1}{r_c} \right)$

- Do the usual KK decomposition:

$$\Psi_{L,R}(x, \phi) = \frac{e^{\frac{3}{2}\sigma}}{\sqrt{r_c}} \sum_n \Psi_n^{L,R}(x) \hat{\phi}_n^{L,R}(\phi) \quad (2)$$

$$\int_{-\pi}^{\pi} d\phi \hat{\phi}_n^{L*}(\phi) \hat{\phi}_m^L(\phi) = \int_{-\pi}^{\pi} d\phi \hat{\phi}_n^{R*}(\phi) \hat{\phi}_m^R(\phi) = \delta_{mn}$$

- The profile of the wavefunction is controlled by $m = ck$. Enters into the order of Bessel fn.

Bulk Fermions II

- The equations are

$$\left(m - \frac{k}{2} + \frac{1}{r_c} \partial_\phi\right) \hat{\phi}_n^R = m_n e^\sigma \hat{\phi}_n^L \quad (3)$$

$$\left(m + \frac{k}{2} - \frac{1}{r_c} \partial_\phi\right) \hat{\phi}_n^L = m_n e^\sigma \hat{\phi}_n^R$$


- The zero modes which we identify as SM fermions $c = \nu \equiv \frac{m}{k}$

$$\hat{\phi}_0^L = N_L^0 e^{kr_c \phi(1/2+\nu)}, \quad N_L^0 = \sqrt{\frac{kr_c(\nu + 1/2)}{e^{2kr_c \pi(\nu+1/2)} - 1}} \quad (4)$$

$$\hat{\phi}_0^R = N_R^0 e^{kr_c \phi(1/2-\nu)}, \quad N_R^0 = \sqrt{\frac{kr_c(1/2 - \nu)}{e^{2kr_c \pi(1/2-\nu)} - 1}}$$

- Since both solutions are Z_2 even at $\phi = 0$, only one of the two is allowed by the Z_2 .
- The RH chiral zero mode lives near the UV (IR) brane if $\nu > 1/2$ ($\nu < 1/2$).
- LH zero mode resides close to UV (IR) brane for $\nu < -1/2$ ($\nu > -1/2$)

Profiles of bulk fermions



The thick red lines are the zero mode wave function of RH chiral fermion.

$$t \equiv e^{-kr_c(\pi-\phi)}$$

Fermion Masses in RS

- The coefficients $c_{L,R}$ control the zero modes i.e peaks at UV or IR
- Localize the Higgs at the IR brane
- Have the zero modes i.e. the SM chiral fermions localize near UV brane
- The overlap after SSB will be very small
- No need to fine tune Yukawa's.
- Quark masses are naturally small.
- If all the fermions both LH doublet and RH singlets are localized near UV then t-quark comes out too light
- t-quark or (t_L, b_L) must not be too far from IR brane

Quark Masses in RS

- The quark masses are given by

$$(M_f^{RS})_{ij} = v_W \frac{\lambda_{5,ij}^f}{kr_c \pi} f_L^0(\pi, c_{f_i}^L) f_R^0(\pi, c_{f_j}^R) \equiv v_W \frac{\lambda_{5,ij}^f}{kr_c \pi} F_L(c_{f_i}^L) F_R(c_{f_j}^R), \quad f = u, d,$$

where the label f denotes up-type or down-type quark species. $v_w = 174$ GeV.

$$f_{L,R}^0(\phi, c_{L,R}) = \sqrt{\frac{kr_c \pi (1 \mp 2c_{L,R})}{e^{kr_c \pi (1 \mp 2c_{L,R})} - 1}} e^{(1/2 \mp c_{L,R}) kr_c \phi}$$

where the upper (lower) sign applies to the LH (RH) zero mode

- The Yukawa couplings λ_{ij} are not necessarily symmetric in i, j
- $f_{L,R}$ shows that the masses are control by values of $c_{L,R}$
- The task is to configurations that fits the CKM matrix.
- Added bonus : both LH and RH quark configurations are given for each solution
- Both **LH and RH rotations** are given for each solution
- In the SM only LH rotations are detectable. $V_{CKM} = V_L^{u\dagger} V_L^d$.

General Configurations

- In general quark mass matrices are not symmetrical in RS. Several configurations found. One example:

$$\begin{aligned}c_Q &= \{0.634, 0.556, 0.256\}, \\c_U &= \{-0.664, -0.536, 0.185\}, \\c_D &= \{-0.641, -0.572, -0.616\}.\end{aligned}\tag{5}$$

- The u and d quark mass matrices (at TeV scale)

$$\langle |M_u| \rangle = \begin{pmatrix} 8.97 \times 10^{-4} & 0.049 & 0.767 \\ 0.010 & 0.554 & 8.69 \\ 0.166 & 9.06 & 142.19 \end{pmatrix}, \quad \langle |M_d| \rangle = \begin{pmatrix} 0.0019 & 0.017 & 0.0044 \\ 0.022 & 0.196 & 0.050 \\ 0.352 & 3.209 & 0.813 \end{pmatrix},$$

where we have used $ke^{-kr_c\pi} = 1.5 \text{ TeV}$.

RS Quark Masses contd

- The CKM matrix elements for the above

$$\begin{aligned} |V_{us}^L| &= 0.16(14), & |V_{ub}^L| &= 0.009(11), & |V_{cb}^L| &= 0.079(74), \\ |V_{us}^R| &= 0.42(24), & |V_{ub}^R| &= 0.12(10), & |V_{cb}^R| &= 0.89(13), \end{aligned} \quad (6)$$

- Note the RH rotations are larger than the LH ones.
- Appears to be true from the numerical searches we found
- How to test it?

Symmetrical Mass Matrices in RS

- Most of the 'constructions' start from conjecture assuming that they are symmetrical
- Put zeros (1 to 3) in appropriate places to fit CKM and the observed mass heirarchies.
- Can RS accomodate these without fine tuning the Yukawa couplings
- Only **ONE** texture zero structures are allowed.
- By construction $U_L = U_R$

All is not well

- The main problem is that the new KK modes will modify EWPT
- The S, T parameters will receive tree level corrections
- It is known that $\rho = 1$ is protected by a custodial $SU(2)$ symmetry
- Promote that to a bulk gauge symmetry.
- Tree level KK gauge effects are suppressed
- The gauge symmetry is now $SU(2)_L \times SU(2)_R \times U(1)_X$
- Take $X = B - L$

Custodial RS model

- Break $SU(2)_R \rightarrow U(1)_R$ by orbifold b.c.

	<i>UV</i>	<i>IR</i>
$\tilde{W}_\mu^{1,2}$	-	+
<i>other gauge fields</i>	+	+

- $U(1)_R \times U(1)_X \rightarrow U(1)_Y$ by vev on **UV** brane. We have a Z' and B_μ

$$Z'_\mu = \frac{g_5 \tilde{W}_\mu^3 - g'_5 \tilde{B}_\mu}{\sqrt{g_5^2 + g'^2_5}}$$

and

$$B_\mu = \frac{g'_5 \tilde{W}_\mu^3 + g_5 \tilde{B}_\mu}{\sqrt{g_5^2 + g'^2_5}}$$

- B_μ is the SM hypercharge gauge boson and broken with $SU(2)_L$ on **IR** brane by Higgs.

Quark Representations

- Zero modes have parity (++)
- Usual assignment

$$\begin{array}{cc} SU(2)_L & SU(2)_R \\ \begin{pmatrix} t_L \\ b_L \end{pmatrix} & \begin{pmatrix} t_R \\ b_R \end{pmatrix} \end{array}$$

because t_R is a zero mode and $SU(2)_R$ is broken on UV


- d_R and t_R must have their own (- +) partners

$$\begin{array}{ccc} SU(2)_L & & SU(2)_R \\ \begin{pmatrix} t_L \\ b_L \end{pmatrix} & \begin{pmatrix} T_R \\ b_R \end{pmatrix} & \begin{pmatrix} t_R \\ B_R \end{pmatrix} \end{array}$$


- They don't affect the quark mass matrix.

FCNC in the Minimal Constrained RS Model

- Besides the direct production of the KK $Z \gtrsim 2.5$ TeV is tree level FCNC
- FCNC $Z - Z_{KK}$ and $Z - Z'_K K$ mixing



- KK-fermion mixings



- Going to the mass basis the unitarity is broken \rightarrow FCNC

$$t \rightarrow Z + jets$$

- The BR is

$$\begin{aligned} Br(t \rightarrow c(u)Z) &= \frac{2}{\cos^2 \theta_W} \left(|Q_Z(t_L) \hat{\kappa}_{tc(u)}^L|^2 + |Q_Z(t_R) \hat{\kappa}_{tc(u)}^R|^2 \right) \left(\frac{1-x_t}{1-y_t} \right)^2 \left(\frac{1+2x_t}{1+2y_t} \right) \\ &= 1.8677 \times \left(|Q_Z(t_L) \hat{\kappa}_{tc(u)}^L|^2 + |Q_Z(t_R) \hat{\kappa}_{tc(u)}^R|^2 \right) \end{aligned}$$

where $x_t = \frac{m_Z^2}{m_t^2}$ and $y_t = \frac{m_W^2}{m_t^2}$ and $Q_Z(f) = T_L^3(f) - Q \sin^2 \theta_w Q_f$

- LH and RH decays are different because $\kappa^R > \kappa^L$ in the config we found

$$\hat{\kappa}_{ab}^u = (U^u)_{a3}^\dagger (\kappa_{Q_3^u}^g + \kappa_{Q_3^u}^f) U_{3b}^u, \quad Q^u = \{u, c, t\}$$

- $BR(t_R \rightarrow Z + c(u)_R) > BR(t_L \rightarrow Z + c(u)_L)$ by ~ 20 .
- The BR is $\sim 10^{-5}$ c.f SM $\sim 10^{-13}$.
- Compare the decays in $t\bar{t}$ vs single tW channels.

Conclusions

- We have found that the RS model can have good quark mass matrices without fine tuning Yukawas
- It can accomodate symmtricall mass matrices if there is only one texture zero and not more
- For asymmetrical conf $U_R > U_L$
- Tree level FCNC best probe in $t \rightarrow Z + jets$
- BR is $\sim 10^{-5}$ makes it very exciting at the LHC
- Predicts RH decays are dominant.